

For Sale

±13.46 AC



Dreher Shoals

Irmo, South Carolina

For more information:

Tombo Milliken

803.744.9852

tombo.milliken@naicolumbia.com

Tom Milliken


803.744.9837

tmilliken@naicolumbia.com

Property Features


- ±13.46 acres located in the Dutch Fork area and fronting Dutch Fork Road in Richland County
- To be used commercially or as a multi-family site
- Market: Columbia Metro
- Submarket: Chapin/Ballentine
- Zoning: Rural, Richland County
- Sales Price: **\$1,204,670 or \$89,500 per acre**

Points of Interest ±13.46 AC



Location

±13.46 AC



Aerial

±13.46 AC



Infrared

±13.46 AC



Topography | 2' Contours
±13.46 AC



Topography | 10' Contours
±13.46 AC



National Wetlands Inventory
±13.46 AC



National Wetlands Inventory

- Estuarine and Marine Deepwater
- Estuarine and Marine Wetland
- Freshwater Emergent Wetland
- Freshwater Forested/Shrub Wetland
- Freshwater Pond
- Lake
- Riverine

Soil Survey
±13.46 AC



Map Unit Description (Brief, Generated)

Richland County, South Carolina

[Minor map unit components are excluded from this report]

Map unit: GeB - Georgeville silt loam, 2 to 6 percent slopes

Component: Georgeville (85%)

The Georgeville component makes up 85 percent of the map unit. Slopes are 2 to 6 percent. This component is on interfluves, piedmonts. The parent material consists of residuum weathered from metavolcanics and/or residuum weathered from metasedimentary rock and/or residuum weathered from slate. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is high. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 2e. This soil does not meet hydric criteria.

Map unit: NaB - Nason silt loam, 2 to 6 percent slopes

Component: Nason (100%)

The Nason component makes up 100 percent of the map unit. Slopes are 2 to 6 percent. This component is on hillslopes on uplands. The parent material consists of clayey residuum weathered from slate. Depth to a root restrictive layer, bedrock, paralithic, is 40 to 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is moderate. Shrink-swell potential is moderate. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 2e. This soil does not meet hydric criteria.

Map unit: NaC - Nason silt loam, 6 to 10 percent slopes

Component: Nason (100%)

The Nason component makes up 100 percent of the map unit. Slopes are 6 to 10 percent. This component is on hillslopes on uplands. The parent material consists of clayey residuum weathered from slate. Depth to a root restrictive layer, bedrock, paralithic, is 40 to 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is moderate. Shrink-swell potential is moderate. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 3e. This soil does not meet hydric criteria.

Map unit: OaB - Orange loam, 0 to 4 percent slopes

Component: Orange (100%)

The Orange component makes up 100 percent of the map unit. Slopes are 0 to 4 percent. This component is on marine terraces on coastal plains. The parent material consists of clayey residuum weathered from slate. Depth to a root restrictive layer, bedrock, lithic, is 40 to 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during January, February, March, December. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 3e. This soil does not meet hydric criteria.