



Ta	Tunica silty clay, 0 to 3 percent slopes, rarely flooded	22.48	1.4%		Illw			625		10.5	55			35	9.5		
Fg	Forestdale silty clay loam, 0 to 0.5 percent slopes	19.91	1.3%		Illw	6.5	50	600				130		35	9		
Sb	Sharkey clay, 0.5 to 3 percent slopes, rarely flooded	19.38	1.2%		IVw	6.5		612.5		10				36.2	9		
Dd	Dubbs very fine sandy loam, 0 to 3 percent slopes, rarely flooded	17.51	1.1%		Ile		90	850		12				40	10		
Tb	Tunica silty clay, 3 to 7 percent slopes	15.57	1.0%		Ille			575		10.5	55			35	9.5		
De	Dubbs very fine sandy loam, 3 to 7 percent slopes, moderately eroded, rarely flooded	14.90	0.9%		IVe		80	750		11				30	8		
Dp	Dundee very fine sandy loam, 3 to 7 percent slopes, rarely flooded	7.06	0.4%		IVe		70	600		8				30	8.5		
W	Water	4.68	0.3%														
Dh	Dundee silt loam, 3 to 7 percent slopes, moderately eroded, rarely flooded	4.60	0.3%		IVe		70	600		8				30	8.5		
Dc	Dubbs silt loam, 0 to 3 percent slopes	3.15	0.2%		Iw		90	850		12				40	10		
Be	Teksob very fine sandy loam, 3 to 7 percent slopes, moderately eroded	2.52	0.2%		IVe		80	750		11				30	8		
Ae	Alligator silty clay, 0.5 to 3 percent slopes	2.11	0.1%		Ille	6.5		550		9		6		30	9	45	
Ac	Alligator clay, 3 to 7 percent slopes	1.25	0.1%		Ille	6.5		550		9		6		30	9	45	
So	Souva silt loam (forestdale)	1.02	0.1%		Illw	6.5	50	600				130		35	9		
Dg	Dundee silt loam, 0 to 3 percent slopes, rarely flooded	0.87	0.1%		Ile	7.5	85	810		13.8				36	9		
Weighted Average					*-		3.4	28.7	387.3		4.5	1.3	13.9	1	22.3	5.4	7.7

*c: Using Capabilities Class Dominant Condition Aggregation Method

* - Non Irr Class weighted average cannot be calculated on the current soils data due to missing data.
Soils data provided by USDA and NRCS.