



County: Bremer County, IA
Location: 25-91N-11W
Township: Franklin
Acres: 113.43



Field borders obtained from Farm Service Agency as of 2008

12/10/2024

FarmWorth, LLC makes no representations or warranties, express or implied, as to the accuracy of any information, data, numerical values, boundaries, or any other information generated through the use of FarmWorth.com. User is solely responsible for independently investigating and determining all information provided through FarmWorth.com prior to use and waives any and all claims against FarmWorth, LLC for any inaccuracies or inconsistencies in the information and/or data.

| Code     | Description                                      | Acres | % of<br>field | IA<br>CSR2 | Overall<br>NCCPI | Soil Class<br>(Irr) | Soil Class (Non-<br>Irr) | Drainage Class             |
|----------|--|-------|---------------|------------|------------------|---------------------|--------------------------|----------------------------|
| 395B     | Marquis loam, 2 to 5 percent slopes              | 32.27 | 28.45 %       | 91         | 90               | -                   | 2                        | Moderately well drained    |
| 399      | Readlyn silt loam, 1 to 3 percent slopes         | 30.54 | 26.92 %       | 91         | 80               | -                   | 1                        | Somewhat poorly<br>drained |
| 84       | Clyde silty clay loam, 0 to 3 percent slopes     | 27.08 | 23.87 %       | 88         | 90               | -                   | 2                        | Poorly drained             |
| 41B 📙    | Sparta loamy sand, 2 to 5 percent slopes         | 7.22  | 6.37 %        | 42         | 60               | 2                   | 4                        | Excessively drained        |
| 175B 🔲   | Dickinson fine sandy loam, 2 to 5 percent slopes | 4.88  | 4.30 %        | 50         | 61               | -                   | 3                        | Well drained               |
| 221      | Klossner muck, 1 to 4 percent slopes             | 4.70  | 4.14 %        | 48         | 89               | -                   | 3                        | Very poorly drained        |
| 471      | Oran loam, 1 to 3 percent slopes                 | 4.14  | 3.65 %        | 74         | 80               | -                   | 1                        | Somewhat poorly<br>drained |
| 198B 🔃   | Floyd loam, 1 to 4 percent slopes                | 2.61  | 2.30 %        | 89         | 88               | -                   | 2                        | Somewhat poorly<br>drained |
| Average: |  |       |               | 83.0       | 83.7             |                     |                          |                            |

IA has updated the CSR values for each county to CSR2. Soils data provided by USDA and NRCS.